



A



B



C



A. The Western Blot analysis of NSE depicted in Figure 2D is presented within the confines of a black rectangle of A. The molecular weight of the NSE protein is determined to be 47kd, and a protein marker is also included on the right side of the image. Notably, the figure retains bands corresponding to markers of 25kd, 35kd, 40kd, 50kd and 70kd. It is important to note that the image has not undergone extensive processing involving high contrast adjustments or multiple exposures, and the primary information is duly marked within the image.

B. The Western Blot analysis of β 3-tubulin depicted in Figure 2D is presented within the confines of a black rectangle of B. The molecular weight of the β 3-tubulin protein is determined to be 55kd, and a protein marker is also included on the left side of the image. Notably, the figure retains bands corresponding to markers of 25kd, 35kd, 40kd, 50kd and 70kd. It is important to note that the image has not undergone extensive processing involving high contrast adjustments or multiple exposures, and the primary information is duly marked within the image.

C. The Western Blot analysis of GAPDH depicted in Figure 2D is presented within the confines of a black rectangle of C. The molecular weight of the GAPDH protein is determined to be 36kd, and a protein marker is also included on the left side of the image. Notably, the figure retains bands corresponding to markers of 35kd, 40kd. It is important to note that the image has not undergone extensive processing involving high contrast adjustments or multiple exposures, and the primary information is duly marked within the image.