```
if (Treatment([diet, oral_antidiabetic], status = PRESCRIBED) and (patient.HbA1c <= 6.5) and
    (not Treatment([diet, metformin], status = CURRENT_OR_PAST, failed = True)) and
    (not Treatment([diet, agi     ], status = CURRENT_OR_PAST, failed = True)) and
    (not Treatment([diet, metformin], status = PRESCRIBED)) and
    (not Treatment([diet, agi     ], status = PRESCRIBED))):
      criticism("Sulfonamides, glinides and glitazones are not recommended at this stage of the disease.
Guideline recommends metformin as first-line treatment.
(Traitement médicamenteux du diabète de type 2)")

if (Treatment([diet, oral_antidiabetic], status = PRESCRIBED) and (patient.A1C <= 6.5) and
    (     Treatment([diet, metformin], status = CURRENT_OR_PAST, failed = True) or
          Treatment([diet, agi     ], status = CURRENT_OR_PAST, failed = True)) and
    (not Treatment([diet, metformin], status = PRESCRIBED)) and
    (not Treatment([diet, agi     ], status = PRESCRIBED))):
      criticism("Sulfonamides, glinides and glitazones are not recommended at this stage of the disease.
Guideline recommends metformin as first-line treatment, and AGI as second-line.
(Traitement médicamenteux du diabète de type 2)")

if (Treatment([diet, oral_antidiabetic], status = PRESCRIBED) and (patient.A1C <= 6.5) and
    (not Treatment([diet, metformin], status = CURRENT_OR_PAST, failed = True)) and
    (not Treatment([diet, agi     ], status = CURRENT_OR_PAST, failed = True)) and
    (not Treatment([diet, metformin], status = PRESCRIBED)) and
    (     Treatment([diet, agi     ], status = PRESCRIBED))):
     criticism("AGI should be prescribed only as second-line treatment.
Guideline recommends metformin as first-line treatment
(Traitement médicamenteux du diabète de type 2)")
```