

Condition	Pneumonia	Diarrhoea	HIV/AIDS	HIV/AIDS	HIV transmission from mother to child	Tuberculosis + HIV	Tuberculosis	Malaria	Malaria
Intervention	Integrated Management of Childhood Illness (IMCI)	Safe water (software)	Highly active anti-retroviral therapy (HAART)	Voluntary Counselling and Testing (VCT)	Single dose Nevirapine (targeted treatment)	Community based DOTs	Community based DOTs	Intermittent treatment of pregnant mothers	Impregnated bed nets
Prevalence (%)	14	12	32.1 * 4.0 **	32.1 * 4.0 **	13.7	44 ***	0.4	9.4	30.9 - 55.4
Burden of disease# Dar es Salaam	5.1	0.9	13.5	13.5	1.4	17.6	3.9	2.9	7.9
Coverage (%) – expert estimates	37	40-50	< 1	< 5	2	100	100	30	12
Data Sources	National Bureau of Statistics, 1999	National Bureau of Statistics, 1999	National AIDS Control Programme, 2000	National AIDS Control Programme, 2000	Kilewo 2001 Kwesigabo 2000	Range 2001	Global TB Programme, WHO, 2002	Mnyika, 2000 Malaria Control Programme, 2002	Drakeley 2000 Wakibara, 1997 Malaria Control Programme, 2002

* = Urban, Iringa; ** = Rural, Mwanza; *** = Percentage of TB-patients with HIV; # = MoH/AMMP data