

Randomized^{1,2}
N=10 Sites (110 patients)

Diabetes Decision Aid
(usual care comparator for Statin Decision Aid)³

(N=5 Sites)

Enrolled patients = 27 DA/38 UC (2 DA & UC)

Median per site: 10 (range 3, 28)

Decline = 65 Median per site: 7 (range 1, 43)

Reason: Unable to reach (48)

Not interested (10)

Other (5)

Evaluable for analysis: N=65 (27 DA)

Missing follow-up: n (DA)

Patient Baseline Survey – 1 (0)

Clinician Survey – 1 (1)

Patient 3 month survey – 13 (7)

Patient 6 month survey – 14 (8)

Statin Decision Aid
(usual care comparator for Diabetes Usual Care)⁴

(N=5 Sites)

Enrolled patients= 26 DA/14 UC (2 DA & UC)

Median per site: 2 (range 2, 28)

Declined = 32 Median per site: 4 (range 1, 17)

Reason: Unable to reach (11)

Not interested (19)

Other (2)

Evaluable for analysis: N=39 (26 DA)

Missing follow-up: n (DA)

Patient Baseline Survey – 3 (1)

Clinician Survey – 7 (4)

Patient 3 month survey – 12 (8)

Patient 6 month survey – 11 (7)

DA = decision aid; UC = usual care

1 – All sites assessed for eligibility agreed to participate and were randomized; none withdrew.

2 – 2 patients removed after aspirin arm of trial was dropped from study.

3 – 2 patients enrolled in both arms were excluded from analysis

4 – 2 patients enrolled in both arms were excluded from analysis (1 is a post randomization exclusion) and 1 UC patient was incorrectly enrolled.