## General practice Hospital 4.0-4.0 -Number of contacts 0.2 0.1 0.1 3.0 2.0-No comorbidity 1.0 0.0 0.0 -8 -12 -11 -10 -9 -12 -11 -10 -9 -4 -3 -2 -1 D -7 -6 -5 -4 -3 -2 Ď -8 -7 -5 -1 -6 3.0 3.0 ł 2.0 2.0 ncidence rate ratios 1.5 1.5 1.2 • • 1.2 1.0 1.0 4.0 4.0 Number of contacts 3.0 3.0 2.0 2.0 Low comorbidity 1.0 1.0 0.0 0.0 -2 -1 D -12 -11 -10 -9 -12 -11 -10 -9 -4 -3 -8 -7 -6 -5 -4 -3 -2 -1 D -8 6 -5 3.0 3.0 ncidence rate ratios 2.0 2.0 1.5 1.5 1.2 1.2 1.0 1.0 4.0 4.0-Number of contacts 3.0 3.0 2.0 2.0 High comorbidity 1.0 1.0 0.0 0.0 -4 -3 -2 -12 -11 -10 -9 -8 -7 -6 -5 -1 Ď -12 -11 -10 -9 -8 -7 -6 -5 -4 -3 -2 -1 Ď 3.0 3.0 Incidence rate ratios 2.0 2.0 1.5-1.5 1.2 1.2 1.0 1.0 Second primary In remission Recurrence

Additional file 5: Number of contacts in men, stratified on healthcare setting and comorbidity burden

Number of contacts are presented as crude rates of mean number of contacts per month. Consultation rate ratios were adjusted for age, comorbidity, educational level, marital status, primary cancer type and time since completion of primary cancer treatment. Patients in remission served as the reference group. Black lines represent 95% confidence intervals.