

Additional Table 3: Treatment modality for gestational diabetes by calendar year

	All	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	<i>P</i> value ¹
	<i>n</i> =	<i>n</i> =	<i>n</i> =	<i>n</i> =	<i>n</i> =	<i>n</i> =	<i>n</i> =	<i>n</i> =	<i>n</i> =	<i>n</i> =	<i>n</i> =	<i>n</i> =	
	30,474	2598	2566	2846	2571	2491	2616	2548	2911	3152	2941	3234	
MNT	18657 (61.2)	1968 (75.8)	1941 (75.6)	2008 (70.6)	1609 (62.6)	1695 (68.0)	1740 (66.5)	1387 (54.4)	1832 (62.9)	1582 (50.2)	1189 (40.4)	1706 (52.8)	0.9995
Antidiabetic oral agents	9942 (32.6)	469 (18.1)	500 (19.5)	705 (24.8)	819 (31.9)	645 (25.9)	739 (28.2)	985 (38.7)	938 (32.2)	1394 (44.2)	1469 (49.9)	1279 (39.5)	0.0003
Insulin therapy	1875 (6.2)	161 (6.2)	125 (4.9)	133 (4.7)	143 (5.6)	151 (6.1)	137 (5.2)	176 (6.9)	141 (4.8)	176 (5.6)	283 (9.6)	249 (7.7)	0.0801

¹Obtained by Mann-Kendall test; the alternative hypothesis states there exists a monotonically increasing trend.