

**Supplementary Table-2: Correlation of major MTBC lineages and site of extrapulmonary infections.**

Lineages	N (%) <a href="#">EPTB[1]</a> (N=1003)	Lymphnode N=626	OR(95% CI):p value	<a href="#">GITB[2]</a> N=168	OR(95% CI):p value	<a href="#">CNSTB[3]</a> N=86	OR(95% CI):p value	Bone & joints N=45	OR(95% CI):p value	<a href="#">UGTB[4]</a> N=41	OR(95% CI):p value	Others N=37
<i>M.bovis</i>	119(11.8)	105(16.8)	<b>5.22(2.23-8.22):&lt;0.001</b>	8(4.8)	<b>0.33(0.085-0.567):0.001</b>	-		3(6.7)	0.52(0.16-1.69): 0.38	3(7.3)	0.57(0.17-1.89): 0.22	-
<i>Delhi/CAS (Indo Oceanic)</i>	275(27.4)	145(23.2)	<b>0.57(0.411-0.734):&lt;0.001</b>	65(38.7)	<b>1.87(1.22-2.53):&lt;0.001</b>	30(34.9)	0.74(0.42-1.06): 0.18	13(28.9)	1.07(0.37-1.79): 0.82	11(26.8)	11.39(2.49-20.28)	11(29.7)
<i>EAI (East African Indian)</i>	158(15.7)	96(15.3)	0.92(0.59-1.24): 0.62	26(15.5)	0.97(0.52-1.42): 0.91	14(16.3)	<b>1.98(0.76-3.19):0.04</b>	8(17.8)	1.16(0.79-1.53): 0.86	6(14.6)	1.66(1.29-2.02): 0.39	8(21.6)
<i>Beijing (East Asian)</i>	43(4.3)	31(4.9)	1.58(0.52-2.64): 0.24	3(1.8)	0.36(0.11-1.18): 0.08	3(3.5)	0.79(0.24-2.61)	2(4.4)	1.04(0.24-4.44)	1(2.4)	0.58(0.07-4.29)	3(8.1)
<a href="#">Euro American[5]</a>	400(39.8)	245(39.1)	0.92(0.79-1.19):0.54	66(39.3)	0.97(0.69-1.36):0.86	37(43)	1.15(0.74-1.80):0.53	18(40)	1.00(0.55-1.85):0.88	20(48.8)	1.48(0.78-2.72):0.24	15(40.5)
<i>-Haarlem</i>	75(7.5)	50(7.9)	1.22(0.61-1.83): 0.50	10(5.9)	0.75(0.23-1.26): 0.51	7(8.1)	1.10(0.20-200): 0.98	3(6.7)	0.88(0.26-2.90)	2(4.9)	0.62(0.15-2.63)	3(8.1)
<i>-LAM</i>	66(6.6)	43(6.9)	1.13(0.54-1.72): 0.73	8(4.8)	0.67(0.16-1.18): 0.38	6(6.9)	1.07(0.44-2.07): 0.88	4(8.9)	1.41(0.48-4.06): 0.74	4(9.7)	1.56(0.54-4.54)	1(2.7)
<i>-Cameroon</i>	49(4.9)	30(4.8)	0.95(0.39-1.51): 0.98	9(5.3)	1.12(0.29-1.96): 0.90	7(8.1)	1.84(0.31-3.38): 0.23	2(4.4)	0.90(0.21-3.83)	1(2.4)	0.47(0.06-3.53): 0.71	-
<i>-Ghana</i>	68(6.8)	41(6.5)	1.94(0.97-2.92): 0.80	13(7.7)	1.18(0.44-1.93): 0.71	5(5.8)	0.84(0.05-1.62): 0.88	3(6.7)	0.98(0.29-3.25)	3(7.3)	1.09(0.32-3.62)	3(8.1)
<i>-Uganda I</i>	41(4.1)	22(3.5)	0.68(0.25-1.12): 0.31	6(3.6)	<b>2.41(0.77-4.06):0.02</b>	5(5.8)	1.51(0.06-2.96): 0.57	3(6.7)	1.73(0.51-5.83): 0.61	5(12.2)	3.57(0.02-7.12): 0.06	-
<a href="#">Others[6]</a>	109(10.87)	53(7.34)	-	20(11.9)	-	9(10.5)	-	7(15.5)	-	5(12.2)	-	8(21.6)

[1] Extrapulmonary tuberculosis

[2] Gastrointestinal tuberculosis

[3] Central nervous system tuberculosis

[4] Urogenital tuberculosis

[5] Lineages other than Delhi/CAS, EAI, Beijing, West African I and II and Bovis together grouped. Sub grouped to analyze 5 major clades

[6] Lineages including Uganda II, West African I and II, X, New-I, S, TUR and undefined lineages