Study or Subgroup	Clindamycin plus quinine Events	Total	Control Events	Total	Weight	Risk Ratio M-H, Fixed, 95% CI
1.2.1 Clindamycin plus quinine vs quinine						
Kremsner 1994	0	39	5	39	45.8%	0.09 [0.01, 1.59]
Metzger 1995	0	40	6	40	54.2%	0.08 [0.00, 1.32]
Subtotal (95% CI)		**79**		**79**	**100.0%**	**0.08 [0.01, 0.63]**
Total events	0		11			
Heterogeneity: Chi² = 0.01, df = 1 (P = 0.94); I² = 0%						
Test for overall effect: Z = 2.42 (P = 0.02)						
1.2.2 Clindamycin plus quinine vs Chloroquine						
Kremsner 1994	0	39	21	39	100.0%	0.02 [0.00, 0.37]
Subtotal (95% CI)		**39**		**39**	**100.0%**	**0.02 [0.00, 0.37]**
Total events	0		21			
Heterogeneity: Not applicable						
Test for overall effect: Z = 2.66 (P = 0.008)						
1.2.3 clindamycin plus quinine vs quinine plus doxycycline						
Metzger 1995	0	40	0	40		Not estimable
Subtotal (95% CI)		**40**		**40**		**Not estimable**
Total events	0		0			
Heterogeneity: Not applicable						
Test for overall effect: Not applicable						
1.2.4 clindamycin plus quinine vs chloroquine plus clindamycin						
Kremsner 1994	0	39	0	39		Not estimable
Subtotal (95% CI)		**39**		**39**		**Not estimable**
Total events	0		0			
Heterogeneity: Not applicable						
Test for overall effect: Not applicable