	Clindamycin plus quinine		Quinine		Risk Ratio		Risk Ratio	
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Fixed, 95% CI	M-H, Fixed, 95% CI	
1.3.1 Clin+quinine vs Quinine: 3-day 12 hourly regimen								
Kremsner 1994	4	39	21	36	48.7%	0.18 [0.07, 0.46]	-	
Metzger 1995	3	40	23	40	51.3%	0.13 [0.04, 0.40]		
Subtotal (95% CI)		79		76	100.0%	0.15 [0.07, 0.32]	•	
Total events	7		44					
Heterogeneity: $Chi^2 = 0.16$, $df = 1 (P = 0.69)$; $I^2 = 0\%$								
Test for overall effect: Z =	5.02 (P < 0.00001)							
1.3.2 Clin+quinine vs Qui	inine: 3-day 8 hourly	regime	n					
Parola 2001	0	57	2	58	100.0%	0.20 [0.01, 4.15]		
Subtotal (95% CI)		57		58	100.0%	0.20 [0.01, 4.15]		
Total events	0		2					
Heterogeneity: Not applica	ıble							
Test for overall effect: Z =	1.04 (P = 0.30)							
1.3.3 Clin+quinine vs Quinine: 7-day 6 hourly regimen								
Pukrittayakamee 2000	0	68	7	68	100.0%	0.07 [0.00, 1.14]		
Subtotal (95% CI)		68		68	100.0%	0.07 [0.00, 1.14]		
Total events	0		7					
Heterogeneity: Not applica	ıble							
Test for overall effect: Z =	1.87 (P = 0.06)							
							0.001 0.1 1 10	1000
						F	avours Clin+Quinine Favours Qu	inine