Additional file 7 Assessment and reporting of risk factors of adverse pregnancy outcomes.

| Study                          | Smoking          | Marital status | Educational<br>level | Height          | Nutrition status | Chronic diseases  |
|--------------------------------|------------------|----------------|----------------------|-----------------|------------------|---|
| Naing, 1988 [26]               | NR               | NR             | NR                   | NR              | NR               | NR  |
| Harinasuta, 1990 [27]          | INA              | INA            | INA                  | INA             | INA              | INA   |
| Nosten, 1993 [28]              | NR‡              | NR‡            | NR‡                  | NR‡             | NR‡              | NR‡   |
| Sowunmi, 1998 [29]             | NR               | NR             | NR                   | NR              | NR               | NR  |
| Bounyasong, 2001 [30]          | NR               | NR             | NR                   | Yes             | NR               | NR  |
| McGready, 2000 [31]            | NR               | NR             | NR                   | NR              | NR               | Women with a history of mental disorder excluded.   |
| McGready, 2001a [32]           | NR               | NR             | NR                   | NR              | NR               | Women with major liver or kidney disease were excluded.   |
| McGready, 2005 [33]            | NR               | NR             | NR                   | NR              | NR               | Women with chronic diseases were excluded from the study.   |
| Adam, 2004a [34]               | NR               | NR             | NR                   | NR              | NR               | NR  |
| Kalilani, 2007 [35]            | NR               | Yes            | Yes                  | NR              | NR               | Women with a history of chronic disease or pregnancy complications were excluded from the study. HIV status was assessed.                               |
| McGready, 2008 [36]            | Yes              | NR             | NR                   | NR              | BMI              | Women with chronic diseases were excluded from the study.   |
| Mutabingwa, 2009 [37]          | NR               | NR             | Yes                  | NR              | NR               | HIV status was assessed. Women with concomitant disease masking assessment of the response to treatment were excluded.                                  |
| Kaye, 2008 [38]                | NR‡              | NR‡            | Yes                  | NR‡             | NR‡              | NR‡   |
| Piola, 2010 [39]               | Recorded but NR. | NR             | NR                   | NR              | NR               | HIV patient was enrolled.   |
| Carmona-Fonseca,<br>2013 [40]  | NR‡              | NR‡            | NR‡                  | NR‡             | NR‡              | NR‡   |
| D'Alessandro, 2016<br>[41, 42] | NR               | NR             | NR                   | NR              | NR               | Women with chronic diseases were excluded. HIV positive patients were enrolled only if she received neither antiretroviral treatment nor cotrimoxazole. |
| Osarfo, 2017 [43]              | NR               | NR             | Yes                  | Yes, but<br>not | NR               | Women with chronic diseases (diabetes, hypertension, renal/cardiac) or HIV were   |
| 0   1   2045 [44]              | 10.1.0           |                |                      | reported        |                  | excluded from the study.  |
| Onyamboko, 2015 [44]           | INA              | INA            | INA                  | INA             | INA              | HIV positive women and women with medical conditions requiring concomitant drug treatment were excluded from the study.                                 |
| Ukah, 2015 [45]                | NR               | NR             | NR                   | NR              | NR               | HIV was assessed and reported.  |
| Iribhogbe, 2017a [46]          | NR‡              | NR‡            | NR‡                  | Yes             | BMI              | Women with known severe underlying diseases were excluded from the study.‡  |
| CTRI/2009/091/001055<br>[47]   | INA              | INA            | INA                  | INA             | INA              | Women with known chronic diseases were excluded from the study.   |
| NCT01054248 [48]               | Yes              | INA            | Yes<br>(literacy)    | Yes             | INA              | Women with a history or known liver diseases or other chronic diseases were excluded.   |
| McGready, 2003a [49]           | NR               | NR             | NR                   | NR              | NR               | Women with chronic diseases were excluded from the study.   |
| Adam, 2012 [50]                | NR               | NR             | NR                   | Yes             | NR               | Women with chronic diseases were excluded.  |
| Onyamboko, 2011 [51]           | NR               | NR             | Yes                  | Yes             | ВМІ              | Women with chronic diseases (diabetes, hypertension, etc.) or HIV were excluded.  |

## Additional file 7 continued.

| Study                       | Smoking                      | Marital status | Educationa<br>level | ll<br>Height | Nutrition status | Chronic diseases   |  |
|-----------------------------|------------------------------|----------------|---------------------|--------------|------------------|--|--|
| McGready,<br>2012 [52]      | Yes                          | NR             | NR                  | NR           | ВМІ              | NR   |  |
| Rijken, 2011<br>[53]        | Yes                          | NR             | NR                  | Yes          | NR               | NR   |  |
| Valea, 2014<br>[54]         | NR                           | Yes            | Yes                 | Yes          | ВМІ              | Patients with chronic medical conditions requiring special care beyond what the study could provide were excluded. |  |
| Juma, 2014<br>[55]          | INA                          | INA            | INA                 | INA          | INA              | Patients with a history of heart disease or arrhythmia were excluded.  |  |
| Mosha, 2014<br>[56]         | No<br>history of<br>smoking. | NR‡            | NR‡                 | Yes          | ВМІ              | No history of chronic disease.   |  |
| Nyunt, 2016<br>[57]         | NR‡                          | NR‡            | NR‡                 | NR           | BMI              | Women with significant other illness including HIV were excluded from the study. ‡                                 |  |
| Mutagonda,<br>2016 [58, 59] | NR‡                          | NR‡            | NR‡                 | NR‡          | NR‡              | NR‡  |  |
| Adam, 2004b<br>[60]         | NR                           | NR             | NR                  | NR           | NR               | NR   |  |
| Adam, 2004c<br>[61]         | NR                           | NR             | NR                  | NR           | NR               | NR   |  |
| Adegnika,<br>2005 [62]      | NR                           | NR             | NR                  | NR           | NR               | NR   |  |
| Adam, 2006<br>[63]          | NR                           | NR             | NR                  | NR           | NR               | NR   |  |
| Ndiaye, 2011<br>[64]        | NR                           | NR             | NR                  | NR           | NR               | Patients with underlying hepatic or renal illness were excluded.   |  |
| Iribhogbe,<br>2017b [65]    | NR‡                          | NR‡            | NR‡                 | Yes          | ВМІ              | Women with known severe underlying diseases were excluded from the study. ‡  |  |
| McGready,<br>1998a [66]     | NR                           | NR             | NR                  | NR           | NR               | NR   |  |
| McGready,<br>1998b [67]     | NR                           | NR             | NR                  | Yes          | NR               | NR   |  |
| McGready,<br>2001b [68]     | NR                           | NR             | NR                  | NR           | NR               | NR   |  |
| Laochan, 2015<br>[69]       | NR‡                          | NR‡            | NR‡                 | NR‡          | NR‡              | NR‡  |  |
| McGready,<br>2002 [70]      | NR                           | NR             | NR                  | NR           | NR               | NR   |  |
| McGready,<br>2003b [71]     | NR                           | NR             | NR                  | NR           | NR               | NR   |  |
| Villegas, 2005<br>[72]      | INA                          | INA            | INA                 | INA          | INA              | INA  |  |
| Rijken, 2008<br>[73]        | NR                           | NR             | NR                  | NR           | NR               | NR   |  |
| Rulisa, 2012<br>[74]        | NR                           | NR             | NR                  | NR           | NR               | Diabetes was reported  |  |
| Kalilani, 2013<br>[75, 76]  | NR                           | Yes            | Yes                 | NR           | NR               | Patients with major illness and HIV-positive patients receiving cotrimoxazole were excluded from the study.        |  |

INA: information not available (conference abstract or registered clinical trial). NR: not reported. ‡: pregnancy outcomes were not reported.