

Additional file 3: Number of districts per malaria intensity within stratification. The West has the most number of district within stratification and the South the lowest. The number of district with high incidence increased across the year mainly in 2013 within stratification.

Stratum	Year	n					
		Low	Lower	Lowest	High	Higher	Highest
East (n= 29)	2010	0	5	10	12	2	0
	2011	2	6	8	9	4	0
	2012	4	4	7	11	2	1
	2013	2	6	8	10	3	0
	2014	6	4	6	9	4	0
West (n= 39)	2010	4	4	14	11	6	0
	2011	4	5	9	17	4	0
	2012	4	5	12	12	6	0
	2013	4	1	15	13	6	0
	2014	5	3	8	17	6	0
South (n=7)	2010	1	3	1	0	2	0
	2011	0	2	2	3	0	0
	2012	0	3	2	1	1	0
	2013	1	0	4	1	1	0
	2014	1	2	2	1	1	0
Fringe (n=16)	2010	2	5	4	3	1	1
	2011	2	4	2	3	5	0
	2012	2	4	3	4	2	1
	2013	1	2	4	5	4	0
	2014	2	3	1	7	2	1
Highlands (n=20)	2010	7	8	2	2	1	0
	2011	5	6	6	2	1	0
	2012	1	4	8	6	1	0
	2013	1	2	9	7	1	0
	2014	4	5	6	2	3	0
Total		65	96	153	168	69	4

n: number of district