

Table S2. Antibodies used for Western blotting

Mouse target proteins	Antigen used to raise antibodies	Type	Species	Dilution	Source	Catalog number	Reference
Cyp2b	CYP2B1	polyclonal	rabbit	1:5000	James Halpert, University of Connecticut	N/A	(1)
Cyp2c	CYP2C11	polyclonal	rabbit	1:10,000	In-house	N/A	(2)
Cyp2d	Cyp2d9	polyclonal	rabbit	1:5000	Masahiko Negishi, NIEHS	N/A	(3)
Cyp2e1	Rat CYP2E1	polyclonal	Rabbit	1:5000	Magnus Ingelman-Sundberg, Karolinska Institute	N/A	(4)
Cyp3a	CYP3A1	polyclonal	rabbit	1:5000	James Halpert, University of Connecticut	N/A	(5)
Cyp4a	CYP4A11 peptide	monoclonal	rabbit	1:2000	Abcam, Cambridge, MA	Ab-140635	(6)
Gapdh	Rabbit GAPDH	monoclonal	mouse	1:10,000	Millipore, Temecula, CA	MAB374	(7)
Nos2	Mouse Nos2	Polyclonal	rabbit	1:2,000	Santa Cruz Biotechnology, Inc, Santa Cruz, CA	Sc-650	(6)

References

1. Duignan DB, Sipes IG, Leonard TB, Halpert JR. Purification and characterization of the dog hepatic cytochrome P-450 isozyme responsible for the metabolism of 2,2',4,4',5,5'-hexachlorobiphenyl. *Arch Biochem Biophys.* 1987;255:290-303.
2. Morgan ET, MacGeoch C, Gustafsson JA. Sexual differentiation of cytochrome P-450 in rat liver. Evidence for a constitutive isozyme as the male-specific 16 alpha-hydroxylase. *Mol Pharmacol.* 1985;27:471-9.
3. Devore K, Harada N, Negishi M. Characterization of testosterone 16 alpha-hydroxylase (I-P-450(16) alpha) induced by phenobarbital in mice. *Biochemistry.* 1985;24:5632-7.
4. Johansson I, Ekstrom G, Scholte B, Puzycki D, Jornvall H, Ingelman-Sundberg M. Ethanol-, fasting-, and acetone-inducible cytochromes P-450 in rat liver: regulation and characteristics of enzymes belonging to the IIB and IIE gene subfamilies. *Biochemistry.* 1988;27:1925-34.

5. Graves PE, Kaminsky LS, Halpert J. Evidence for functional and structural multiplicity of pregnenolone-16 alpha-carbonitrile-inducible cytochrome P-450 isozymes in rat liver microsomes. *Biochemistry*. 1987;26:3887-94.
6. Mimche SM, Nyagode BA, Merrell MD, Lee CM, Prasanphanich NS, Cummings RD, et al. Hepatic cytochrome P450s, phase II enzymes and nuclear receptors are downregulated in a Th2 environment during *Schistosoma mansoni* infection. *Drug Metab Dispos*. 2014;42:134-40.
7. Lee CM, Tripathi S, Morgan ET. Nitric oxide-regulated proteolysis of human CYP2B6 via the ubiquitin-proteasome system. *Free Radic Biol Med*. 2017;108:478-86.