APPENDIX 2: Study outcomes stratified by rural/urban residence

Table A2.1 Use of 4+ ANC among most recent births

	4+ ANC (all women)		4+ ANC (rural)		4+ ANC (urban)	
	Estimate [95% CI]	p-value	Estimate [95% CI]	p-value	Estimate [95% CI]	p-value
Pre-policy	62.3%		56.1%		84.0%	
starting level	[57.4%,67.1%]		[50.1%,62.1%]		[78.7%,89.2%]	
Pre-policy half-	-1.2%	0.000	-0.9%	0.096	-2.3%	<0.001
yearly trend	[-2.2%,-0.3%]	0.009	[-1.9%,0.2%]	0.096	[-3.1%,-1.6%]	\U.UU1
Immediate	+0.3%	0.965	-2.1%	0.720	+11.4%	0.019
change in level	[-11.8%,12.3%]	0.905	[-14.3%,10.2%]	0.730	[2.0%,20.7%]	0.019
Immediate	+2.4%	0.001	+1.8%	0.010	+2.9%	د0 001
change in slope	[1.1%,3.6%]	0.001	[0.5%,3.1%]	0.010	[2.0%,3.9%]	<0.001
Post-policy half-	+1.1%	0.002	+0.9%	0.000	+0.6%	0.061
yearly trend	[0.4%,1.8%]	0.003	[0.2%,1.5%]	0.009	[0.0%,1.2%]	0.061

Table A2.2: Early ANC initiation among users of 1+ ANC

	Early ANC (all women)		Early ANC (rural)		Early ANC (urban)	
	Estimate [95% CI]	p-value	Estimate [95% CI]	p-value	Estimate [95% CI]	p-value
Pre-policy	14.0%		10.7%		23.0%	
starting level	[10.2%,17.9%]		[6.9%,14.6%]		[18.4%,27.6%]	
Pre-policy half-	-0.3%	0.355	0.0%	0.000	-1.0%	0.010
yearly trend	[-0.9%,0.3%]	0.355	[-0.5%,0.6%]	0.899	[-1.8%,-0.2%]	0.019
Immediate	+2.6%	0.202	+0.1%	0.057	+6.0%	0.105
change in level	[-2.5%,7.6%]	0.303	[-4.6%,4.9%]	0.957	[-3.0%,15.0%]	0.185
Immediate	+1.0%	0.000	+0.4%	0.215	+2.1%	0.001
change in slope	[0.3%,1.7%]	0.008	[-0.2%,1.1%]	0.215	[0.9%,3.2%]	0.001
Post-policy half-	+0.7%	<0.001	+0.4%	0.001	+1.1%	0.002
yearly trend	[0.5%,0.9%]	<0.001	[0.2%,0.7%]	0.001	[0.4%,1.7%]	0.002

Table A2.3: Use of ANC from a public sector health facility among users of 1+ ANC

	Any public facility (all women)		Any public facility (rural)		Any public facility (urban)	
	Estimate [95% CI]	p-value	Estimate [95% CI]	p-value	Estimate [95% CI]	p-value
Pre-policy	66.0%		65.1%		64.8%	
starting level	[59.6%,72.4%]		[56.5%,73.6%]		[58.4%,71.2%]	
Pre-policy half-	+1.0%	0.044	+1.2%	0.050	+1.2%	0.031
yearly trend	[0.0%,2.0%]	0.044	[0.0%,2.5%]	0.050	[0.1%,2.4%]	0.031
Immediate	+3.0%	0.400	+2.6%	0.623	-4.1%	0.507
change in level	[-6.1%,12.1%]	0.499	[-8.1%,13.3%]	0.023	[-16.7%,8.5%]	0.507
Immediate	-1.0%	0.042	-0.9%	0.120	-1.4%	0.022
change in slope	[-2.0%,0.0%]	0.042	[-2.2%,0.3%]	0.129	[-2.7%,-0.2%]	0.023
Post-policy half-	0.0%	0.076	+0.3%	0.007	-0.2%	0.502
yearly trend	[-0.2%,0.2%]	0.976	[0.1%,0.5%]	0.007	[-0.8%,0.4%]	0.502

Table A2.4: Use of primary care facility among users of public facility-based ANC

	Primary care facility (all women)		Primary care facility		Primary care faciltiy	
	•		(rural)		(urban)	
	Estimate	p-value	Estimate	p-value	Estimate	p-value
	[95% CI]		[95% CI]		[95% CI]	
Pre-policy	64.5%		69.4%		49.0%	
starting level	[59.2%,69.8%]		[63.7%,75.1%]		[33.3%,64.7%]	
Pre-policy half-	+0.4%	0.256	+0.4%	0 207	+0.1%	0.017
yearly trend	[-0.5%,1.4%]	0.356	[-0.5%,1.2%]	0.387	[-2.2%,2.4%]	0.917
Immediate	-4.7%	0.225	-3.1%	0.276	-13.0%	0.202
change in level	[-14.6%, 5.2%]	0.335	[-10.2,4.0%]	0.376	[-36.4%,10.4%]	0.263
Immediate	-0.7%	0.103	-0.2%	0.645	+0.3%	0.800
change in slope	[-1.7%,0.4%]	0.193	[-1.0%,0.6%]	0.645	[-2.3%,2.9%]	0.800
Post-policy half-	-0.2%	0.247	+0.2%	0.042	+0.4%	0.440
yearly trend	[-0.7%,0.2%]	0.247	[0.0%,0.3%]	0.042	[-0.7%,1.6%]	0.440

Table A2.5: Received good content of care among users of public facility-based ANC

	Received all 6 routine ANC components		Received all 6 routine ANC components		Received all 6 routine ANC components	
	(all women)		(rural)		(urban)	
	Estimate	p-value	Estimate	p-value	Estimate	p-value
	[95% CI]		[95% CI]		[95% CI]	
Pre-policy	9.4%		7.5%		13.4%	
starting level	[4.7%,14.2%]		[3.0%,12.0%]		[6.4%,20.4%]	
Pre-policy half-	+0.4%	0.402	+0.4%	0.353	+0.7%	U 33E
yearly trend	[-0.6%,1.4%]	0.402	[-0.5%,1.3%]	0.353	[-0.8%,2.3%]	0.335
Immediate	+4.9%	0.212	+3.9%	0.250	+8.0%	0.347
change in level	[-4.9%,14.6%]	0.313	[-4.6%,12.5%]	0.350	[-9.2%,25.3%]	0.547
Immediate	+1.1%	0.087	+0.8%	0.162	+0.7%	0.337
change in slope	[-0.2%,2.3%]	0.067	[-0.3%,1.9%]	0.102	[-0.8%,2.1%]	0.557
Post-policy half-	+1.5%	<0.001	+1.2%	<0.001	+1.4%	<0.001
yearly trend	[1.0%,2.0%]	<0.001	[0.7%,1.7%]	<0.001	[1.0%,1.9%]	<0.001

APPENDIX 2: Study outcomes stratified by rural/urban residence

Table A2.6: Summary of the impact of the 10/20 policy on ANC

	Immediate change in	Immediate change in				
	level	slope				
(2) 4+ ANC (most recent births)						
All women	none	increased				
Rural	none	increased				
Urban	increased	increased				
(3) Early ANC (users of 1+ ANC)						
All women	none	increased				
Rural	none	none				
Urban	none	increased				
(4) Public facility-based ANC (users of 1+ ANC)						
All women	none	decreased				
Rural	none	none				
Urban	none	decreased				
(5) Public primary care facility (users of any public facility-based care)						
All women	none	none				
Rural	none	none				
Urban	none	none				
(6) Received good content of ANC (users of any public facility-based care)						
All women	none	none				
Rural	none	none				
Urban none none						
increased: increasing effect or trend, p<0.05						
decreased: decreasing effect or trend, p<0.05						
none: no effect, p>0.10						