Estimating loss in quality of life associated with asthma-related crisis events (ESQUARE): a cohort, observational study Additional file 4

Table 1: Week 8 statistics for each quality of life questionnaires

Item	N	Mean	SD	Range	Response rates	Floor effects	Ceiling effects
EQ-5D-5L (utility)	65	0.725	0.292	-0.215to 1.00	53.7%	0.00%	21.5%
<b>EQ-5D VAS score</b>	65	68.06	21.91	5.00 to 100.00	53.7%	0.00%	0.00%
AQLQ overall score	65	4.48	1.50	1.47 to 7.00	53.7%	0.00%	0.00%
<b>AQLQ Symptoms</b>	66	3.64	2.22	0.00 to 7.00	54.5%	0.00%	0.00%
score							
<b>AQLQ Activity</b>	66	3.68	2.13	0.00 to 7.00	54.5%	0.00%	0.00%
score							
<b>AQLQ Emotional</b>	66	3.72	2.39	0.00 to 7.00	54.5%	0.00%	1.50%
score							
AQLQ	66	3.91	2.33	0.00 to 7.00	54.5%	0.00%	
Environmental							1.50%
score							
AQL-5D (utility)	64	0.737	0.176	0.450 to 1.00	52.9%	0.00%	7.80%
TTO (utility)	80	0.787	0.295	0.000 to 1.00	71.4%*	0.00%	51.3%

<sup>\*</sup>The response rate is based on the denominator being 112 due to only the participants based at the Norwich hospital site (NNUH) being asked the TTO questions. All of the other response rates for the PROMS were based on the denominator being 121 as this was the total number recruited across all hospital sites where each participant was asked to complete PROM questionnaires.

Ranges for PROMs: EQ-5D-5L (-0.281 to 1); EQ-5D VAS (0 to 100); AQLQ (0 to 7); AQL-5D (0 to 1); TTO (0 to 1).