Kamradt et al. (2018) Sustainable reduction of antibiotics-induced antimicrobial resistance (ARena) in German ambulatory care: study protocol of a cluster randomized trial – Additional file 2

Additional file 2

 Table 1 Eligible medical specialist groups for participation in the ARena study

Medical specialist group	German equivalent (Facharztgruppe-FG)
General practitioner (GP)	Allgemeinmediziner, praktischer Arzt, hausärztlich tätige Internist (FG 01, 02, 03)
Internist	Internist (FG 23)
Gynecologist	Gynäkologe (FG 15)
Ear, nose and throat (ENT) specialist	Hals-Nasen-Ohren (HNO) Arzt (FG 19)
Urologist	Urologe (FG 67)
Pulmonary specialist	Pneumologe (FG 30)
Pediatrician	Kinderarzt (FG 34, 46)

 Table 2 Description of the recruitment of participants in the ARena study

 Paceruitment strategy

	Recruitment strategy
General study participation /	In total, 221 ambulatory practices (302 physicians, respectively) in the fourteen
outcome evaluation	practice networks will be invited to participate in the study (arm A: 68
	practices / 102 physicians; arm B: 56 practices / 91 physicians; arm C 68
	practices / 112 physicians). All practices will be included in the claims-data
	based outcome evaluation. The comparison group will be selected randomly
	from ambulatory practices outside of the participating networks. Additionally,
	research-practices of the Dept. of General Practices and Health Services
	Research will be invited for participation in the patient survey pilot study.
Patient survey	Per practice (in study arm B) 60 patients aged 18 years or above and insured at
	the AOK health insurance in Bavaria with index diseases will be asked by their
	physician to participate (anonymously and voluntarily) in the patients survey
	by filling in the questionnaire (main study). The pilot study is conducted in
	practices which do not take part in the ARena main study (10-12 practices, 10-
	15 patients per practice). In the pilot study one or two patients per practice are
	asked by their doctor, in addition to filling in the questionnaire, to participate
	in a short telephone interview conducted by a researcher. All patients receive
	an information leaflet concerning the aim and the background of the survey,
	including information on data protection.
Process evaluation	Physicians in all three intervention arms and non-physician health
	professionals of the practice team (medical assistants) in intervention arm B,
	who were contacted for general study participation, already received
	information about the process evaluation survey through their information
	leaflet.
	Selection of potential interview partners within the sample of physicians and
	medical assistants willing to participate in the intervention study, and
	stakeholders will be based on the strategy of purposive sampling. This
	sampling technique helps to identify individuals that are especially
	knowledgeable about and/or experienced in the phenomenon of interest.
	Every participating practice network is asked to name a GP, who is involved in
	the ARena-interventions and who is interested to take part in the focus group.