			II ROUND DELPHI					
COLOR CODE	Surgical disease/ conditions	Decision I ROUND	Likert scale 4 (near totally agree)	%	Likert scale 5 (totally agree)	0/0	Likert scale 5/4+5	Decision
RED CODE- immediate surgery	blunt and penetrating trauma	INCLUDED	-		-	-		-
	postoperative hemorrhage	INCLUDED	-	-	-	-		-
	post-partum hemorrhage	INCLUDED	-	-	-	-		-
	ruptured aneurysm	INCLUDED	-	-	-	-		-
	Tension pneumothorax	INCLUDED	-	-	-	-		-
ORANGE CODE-surgery within 1 hrs	acute mesenteric ischemia	INCLUDED	-	-	-	-		-

	acute limbs ischemia	INCLUDED	-	-	-	-		-
	graft thrombosis	INCLUDED	-	-	-	-		-
	testicular torsion	INCLUDED	-	-	-	-		-
	retained placenta with acute bleeding	INCLUDED	-	-	-	-		-
Patients with septic shock	GI perforation	INCLUDED	-	-	-	-		-
	Infected necrotizing haemorrhage	II ROUND	8/51	15.7	22/51	43	58.7	Not included, but why?
	pancreatitis							Better definition ? III
	Urolithiasis complicated by hydronephrosi s	II ROUND	14/51	27.5	22/51	43	70.5	Not included, better definition?
	Fournier's gangrene	INCLUDED	-	-	-	-		

	toxic megacolon	INCLUDED	-	-	-	-		
	anastomotic fis tula	INCLUDED	-	-	-	-		
	necrotizing fascitis	INCLUDED	-	-	-	-		
	ruptured tubo- ovarian abscess	II ROUND	12/51	23.5	24/51	47	70.5	Not included, why if patient is in septic shock?
								Better definition III
	Incomplete abortion	II ROUND	13/51	25.5	24/51	47	72.5	III
	Perianal abscess	II ROUND	12/51	23.5	26/51	51	74.5	III
Other conditions	pneumothorax	INCLUDED		-	_			
	heart lung transplants	INCLUDED	-	-	-			

YELLOW- surgery within 3/6	Gangrenous cholecystitis	II ROUND (added)	15/51	29.4	26/51	56. 9	86.3	INCLUD ED
hrs	spinal cord compression with cauda equina syndrome	INCLUDED	-	-	-			
Patients with sepsis associated	Fournier's gangrene	II ROUND	6/51	11.8	35/51	68. 6	80.4	INCLUD ED
with:	toxic megacolon	II ROUND	15/51	29.4	31/51	60. 8	90.2	INCLUD ED
	anastomotic fis tula with local peritonitis	II ROUND	18/51	35.3	17/51	33	68.3	III
	necrotizing fascitis	II ROUND	8/51	15.7	33/51	64. 7	80.4	INCLUD ED
	GI perforation	II ROUND	6/51	11.8	39/51	76. 5	88.3	INCLUD ED
	Ruptured tubo-ovarian abscess	II ROUND	16/51	31.4	26/51	51	82.4	INCLUD ED

	complicated appendicitis	II ROUND	16/51	31.4	21/51	41.	72.6	Better definiti on?III
	Complicated diverticulitis	II ROUND	16/51	31.4	17/51	33.	64.7	Better definiti on?III
	Urolithiasis with sepsis	II ROUND	13/51	25.5	17/51	33.	58.8	Better definiti on?III
	Incomplete abortion	II ROUND	15/51	29.4	27/51	52. 9	82.3	INCLUD ED
	Perianal abscess with sepsis	II ROUND	17/51	33.3	20/51	39. 2	72.5	Better definiti on?III
Other conditions	Intraperitoneal bladder rupture	II ROUND	18/51	35.3	19/51	37. 3	72.3	Why? Patient in sepsis?
	compartment syndrome (any districts)	II ROUND	8/51	15.7	34/51	66.	82.4	INCLUD ED

	foreign body with GUT obstruction (including endoscopy)	II ROUND	18/51	35.3	15/51	29. 4	64.7	Better definiti on?
	liver transplant	INCLUDED	-	-	-	-		-
	Embolization of post-traumatic pseudoaneurys m	II ROUND added	16/51	31.4	21/51	41.	72.6	III
GREEN-surgery within 12 hrs	bowel/ intestinal obstruction	II ROUND	15/51	29.4	19/51	37.	66.7	? III
	Perianal abscess	II ROUND	12/51	23.5	21/51	41.	64.7	? III
	Cholecystitis	II ROUND	13/51	25.5	18/51	35. 3	60.8	? III

Appendicitis with systemic signs of infection	II ROUND	11/51	21.6	32/51	62. 7	84.3	INCLUD ED
Thoracic empyema	INCLUDED	-	-	-			-
incarcerated hernia with obstruction	II ROUND	9/51	17.6	28/51	54. 9	72.5	? III
Urinary fistula	II ROUND	9/51	17.6	11/51	21.	39.2	Cancell ed
Hydronephrosi s (stent placement)	II ROUND	13/51	25.5	16/51	31.	56.9	?III
Urolithiasis with acute kidney injury	II ROUND	13/51	25.5	16/51	31.	56.9	? III
kidney and pancreatic transplant	II ROUND	13/51	25.5	21/51	41.	66.7	? III

	foreign body without obstruction	II ROUND	9/51	17.6	12/51	23. 5	41.1	Cancell ed
BLUE CODE- surgery within 24-48hrs	2nd look laparotomy	INCLUDED	-	-	-	-		-
	cholecystectom y after ERCP for stones migration or recurrent biliary colic	INCLUDED	-	-	-	-		-
	Uncomplicated appendicitis	II ROUND	7/51	13.7	31/51	60.	74.5	? III
	Amputation for osteomyelitis	II ROUND	6/51	11.8	30/51	58. 8	70.6	? III
	A-V fistulae for hemodialysis	II ROUND	8/51	15.7	28/51	54. 9	70.6	? III

Symptomatic carotid artery stenosis, TIA, STROKE in evolution	II ROUND	8/51	15.7	19/51	37.	53	? III
Symptomatic AAA after medical treatment failure	II ROUND	15/51	29.4	21/51	41.	70.6	? III
Symptomatic aortic dissection type B	II ROUND	9/51	17.6	21/51	41.	58.8	? III
Bones fracture	II ROUND	13/51	25.5	22/51	43. 1	68.6	? III
pelvic trauma fixation	II ROUND	13/51	25.5	27/51	52. 9	78.4	INCLUD ED
Maxillofacial fractures	II ROUND	12/51	23.5	25/51	49	72.5	III
Urolithiasis after medical treatment failure	II ROUND	13/51	25.5	19/51	37. 3	62.8	? III

ORGANIZATIV E NEED	Elective postponed interventions	II ROUND	12/51	23.5	28/51	54. 9	82.9	INCLUD ED
	diagnostic biopsy/ laparoscopy	II ROUND	9/51	17.6	30/51	58. 8	76.4	INCLUD ED