

## PRENATAL STRESS AND EXTERNALIZING BEHAVIOR

### Additional file Tables

Table S1: Pregnancy perceived stress with absent vs. present medical risk factors during pregnancy (Welch's test)

	PSQ Risk absent		PSQ Risk present		t	Welch's df	p
	M	SD	M	SD			
	Pregnancy complications t1	38.844	20.03	42.425			
Pregnancy complications t2	37.855	19.14	40.873	19.83	-.75	66.09	.467
Pregnancy complications t3	37.596	19.93	43.536	18.50	-1.55	89.48	.125
Alcohol consumption t1	39.596	19.71	48.148	17.56	-1.15	6.07	.293
Alcohol consumption t2	38.882	19.52	38.334	3.92	.15	4.69	.879
Alcohol consumption t3	40.130	19.70	34.444	16.46	.75	4.89	.490
Hospitalization t1	40.50	19.95	37.89	18.64	.42	11.76	.683
Hospitalization t2	38.91	19.46	38.63	19.22	.05	18.26	.960
Hospitalization t3	39.13	19.94	41.97	18.47	-.68	50.14	.503

Note: t1 = 1<sup>st</sup> pregnancy trimester, t2 = 2<sup>nd</sup> pregnancy trimester, t3 = 3<sup>rd</sup> pregnancy trimester

Table S2: Pregnancy cortisol with absent vs. present medical risk factors during pregnancy (Welch's test)

	Cortisol Risk absent		Cortisol Risk present		t	Welch's df	p
	M	SD	M	SD			
	Pregnancy complications t1	2.90	1.22	2.88			
Pregnancy complications t2	2.81	1.12	3.02	1.05	-.87	72.24	.389
Pregnancy complications t3	2.71	1.19	3.08	.97	-1.70	96.86	.092
Alcohol consumption t1	2.89	1.15	2.86	.95	.02	4.90	.988
Alcohol consumption t2	2.89	1.13	2.51	.86	.61	1.22	.637
Alcohol consumption t3	2.85	1.13	2.97	1.01	-.19	2.31	.812
Hospitalization t1	2.92	1.12	2.94	1.13	-.07	11.43	.942
Hospitalization t2	2.89	1.14	2.86	1.06	.09	17.29	.927
Hospitalization t3	2.82	1.17	2.95	.98	-.55	52.95	.581

Note: t1 = 1<sup>st</sup> pregnancy trimester, t2 = 2<sup>nd</sup> pregnancy trimester, t3 = 3<sup>rd</sup> pregnancy trimester