

'Preclinical care' data set (n = 2.973; 2004-2010)

ROSC = 1.302 (44%)
Admission to Hospital = 1.040 (35%)

'Postresuscitation care' data set (n = 711; 2004-2010)

Location of cardiac arrest:

Home:	363	(51%)	Nursing home:	43	(6%)
Work place:	26	(4%)	Doctors office:	38	(5%)
Public street:	102	(14%)	Others:	136	(19%)

Excluded:
n = 127

Cardiac (n = 466)

Witnessed: 358 (77%)
Bystander CPR: 86 (18%)
VF: 226 (48%)
Non-VF: 240 (52%)

Non-cardiac (n = 118)

Witnessed: 82 (69%)
Bystander CPR: 16 (14%)
VF: 16 (14%)
Non-VF: 102 (86%)

VF (n = 226)

Treatment:

PCI: 47 (21%)
PCI+MTH: 53 (23%)
MTH: 37 (16%)

Outcome:

24 h Survival 183 (81%)
Hospital Disc. 112 (50%)
CPC 1+2 76 (34%)
CPC 3+4 11 (5%)
Dead 110 (49%)

non VF (n = 240)

Treatment:

PCI: 29 (12%)
PCI+MTH: 17 (7%)
MTH: 42 (18%)

Outcome:

24 h Survival 144 (60%)
Hospital Disc. 68 (28%)
CPC 1+2 44 (18%)
CPC 3+4 10 (4%)
Dead 168 (70%)

VF (n = 16)

Treatment:

PCI: 1 (6%)
PCI+MTH: 1 (6%)
MTH: 2 (13%)

Outcome:

24 h Survival 12 (75%)
Hospital Disc. 7 (44%)
CPC 1+2 5 (31%)
CPC 3+4 2 (13%)
Dead 9 (56%)

non VF (n = 102)

Treatment:

PCI: 4 (4%)
PCI+MTH: 2 (2%)
MTH: 25 (25%)

Outcome:

24 h Survival 60 (59%)
Hospital Disc. 17 (17%)
CPC 1+2 12 (12%)
CPC3+4 5 (5%)
Dead 77 (75%)