

Supplementary Table 1. Frequency of the sub-species of the isolated bacteria

Species	Frequency (%)
<i>Escherichia coli</i>	377 (37.3)
<i>Klebsiella</i> species	200 (19.8)
<i>Klebsiella pneumoniae</i>	172
<i>Klebsiella oxytoca</i>	26
<i>Klebsiella aerogenes</i>	2
<i>Staphylococcus</i> species	83 (8.2)
<i>Staphylococcus aureus</i>	55
<i>Staphylococcus epidermidis</i>	17
<i>Staphylococcus capitis</i>	7
Other Coagulase-negative <i>Staphylococci</i> ¹	4
<i>Streptococcus</i> species	63 (6.2)
<i>Streptococcus anginosus</i>	17
<i>Streptococcus pneumoniae</i>	15
<i>Streptococcus agalactiae</i>	11
Other <i>Streptococci</i> ²	20
<i>Enterococcus</i> species ³	57 (5.6)
<i>Enterobacter</i> species ⁴	53 (5.2)
<i>Pseudomonas aeruginosa</i>	51 (5.0)
<i>Citrobacter freundii</i>	21 (2.1)
<i>Acinetobacter baumannii</i>	20 (2.0)
<i>Clostridium</i>	14 (1.4)
Etc. ⁵	71 (7.0)

Data are presented as n (%).

¹Other Coagulase-negative *Staphylococci* included *Stap. haemolyticus*, *Stap. lugdunensis*, *Stap. saccharolyticus*.

²Other *Streptococci* included *Strep. constellatus*, *Strep. dysgalactiae*, *Strep. pyogenes*.

³*Enterococcus* species contained *E. faecium*, *E. faecalis*, *E. casseliflavus*, *E. gallinarum*, *E. raffinosus*.

⁴*Enterobacter* species contained *E. faecium*, *E. cloacae*, *E. aerogenes*, *E. avium*, *E. kobei*, *E. sakazakii*.

⁵Et cetera included *Bacillus* species, *Bacteroides* species, *Campylobacter* species, *Chryseobacterium indologenes*, *Eggerthella lenta*, *Flayonifractor plautii*, *Fusobacterium* species, *Haemophilus influenzae*, *Kocuria* species, *Morganella morganii*, *Peptostreptococcus* species, *Proteus mirabilis*, *Vibrio vulnificus*.