

Supplementary Table 2 Patient characteristics of the different cohorts included in the study.

	Stockholm 2	Stockholm 3	Van de Vijver	Uppsala	Karolinska
	n (%)				
Tumour size					
< 20 mm	37 (43.0)	580 (77.2)	155 (53.0)	127 (25.4)	89 (62.2)
>20 mm	49 (57.0)	171 (22.8)	140 (47.0)	124 (24.8)	54 (37.8)
Lymph node status					
-	6 (7.0)	768 (100)	151 (51.2)	158 (65.3)	87 (60.8)
+	80 (93.0)	0 (0)	144 (48.8)	84 (34.7)	56 (39.2)
Grade	N/A				
1		122 (18.6)	75 (25.4)	67 (26.9)	27 (18.9)
2		382 (58.2)	101 (34.2)	128 (51.4)	57 (39.9)
3		152 (23.2)	119 (40.3)	54 (21.7)	59 (41.3)
ER					
-	18 (20.9)	159 (21.3)	69 (23.4)	34 (13.8)	25 (17.5)
+	68 (79.1)	589 (78.7)	226 (76.6)	213 (88.7)	118 (82.5)
PgR	N/A		N/A		N/A
-		327 (47.6)		61 (24.3)	
+		360 (52.4)		190 (75.7)	
HER2			N/A	N/A	N/A
-	71 (77.2)	634 (88.7)			
+	21 (22.8)	81 (11.3)			