

	Randomized Trials			Open-label Extension Trials <sup>b</sup>	Patients With ≥1 Dose of TCZ <sup>c</sup> (Randomized + Open-label Extension Trials)
1	N = 23 (MTX ≥8 weeks)	• Weekly MTX + single-dose 10 mg/kg TCZ	8 wks	8 mg/kg TCZ ± MTX/DMARDs 5 y	n = 2263
2	N = 673 (6-mos MTX free)	• MTX • 8 mg/kg TCZ	24 wks		
3	N = 499 (TNF-IR)	• Placebo • 4 mg/kg TCZ + MTX • 8 mg/kg TCZ + MTX	24 wks		
4	N = 1220 (DMARD-IR)	• Placebo • 8 mg/kg TCZ + DMARDs	24 wks		
5	N = 623 (DMARD-IR)	• Placebo • 4 mg/kg TCZ + MTX • 8 mg/kg TCZ + MTX	24 wks	8 mg/kg TCZ + MTX 5 y	n = 597
6	N = 1196 (DMARD-IR)	• Placebo • 4 mg/kg TCZ + MTX • 8 mg/kg TCZ + MTX	2 yrs	8 mg/kg TCZ + MTX 3-5 y	n = 1149
<b><u>ALL-CONTROL POPULATION (n = 4199)<sup>a</sup></u></b> 774 patients initially randomized to 4 mg/kg TCZ 1870 patients initially randomized to 8 mg/kg TCZ 1555 patients initially randomized to control (MTX/DMARDs or placebo)			<b><u>ALL-EXPOSED POPULATION (n = 4009)</u></b> 774 patients initially randomized to 4 mg/kg TCZ 1870 patients initially randomized to 8 mg/kg TCZ 1365 patients initially randomized to control (MTX/DMARDs or placebo)		