

Identification of progressors by C1M

Table 2: Univariate correlation between clinical markers and baseline serum C1M.

	Baseline (all)			$\Delta 24$ weeks (PBO)			$\Delta 24$ weeks (PBO w/o escape)			$\Delta 52$ weeks (PBO w/o escape)		
	ρ	P	n	ρ	P	n	ρ	P	n	ρ	P	n
Age	-0.06	ns	585	-	-	-	-	-	-	-	-	-
BMI	0.01	ns	578	-	-	-	-	-	-	-	-	-
Disease duration	-0.01	ns	584	-	-	-	-	-	-	-	-	-
Serum CRP	0.80	<0.0001	584	-0.31	0.001	112	-0.31	0.0023	93	-0.36	0.0007	83
VAS pain	0.21	<0.0001	578	-0.08	ns	110	-0.05	ns	91	-0.00	ns	83
HAQ	0.21	<0.0001	535	-0.17	ns	98	-0.15	ns	82	-0.11	ns	76
DAS28	0.24	<0.0001	573	0.09	ns	108	0.15	ns	89	0.11	ns	77
JSN	0.12	0.0056	556	0.08	ns	153	0.16	ns	90	0.21	0.055	84
mTSS	0.14	0.0006	556	0.19	0.017	153	0.28	0.0084	90	0.34	0.0017	84
ERN	0.15	0.0003	556	0.19	0.019	153	0.13	ns	90	0.21	0.054	84

BMI: Body mass index. CRP: C-reactive protein. VAS pain: Visual analog scale of pain. HAQ: Health assessment questionnaire. DAS28: Disease activity score 28. JSN: Joint space narrowing. mTSS: modified total Sharp score. PBO: placebo group. Spearman rank correlation coefficient ρ , the P values and n are provided.