## Additional File 9

Endo-Lysosomal Proteins and Ubiquitin CSF Concentrations in Alzheimer's and Parkinson's Disease

Simon Sjödin<sup>1,2</sup>, Gunnar Brinkmalm<sup>1,2</sup>, Annika Öhrfelt<sup>1,2</sup>, Lucilla Parnetti<sup>3</sup>, Silvia Paciotti<sup>4</sup>, Oskar Hansson<sup>5,6</sup>, John Hardy<sup>7</sup>, Kaj Blennow<sup>1,2</sup>, Henrik Zetterberg<sup>1,2,7,8</sup>, Ann Brinkmalm<sup>1,2</sup>

<sup>1</sup>Department of Psychiatry and Neurochemistry, Institute of Neuroscience and Physiology, the Sahlgrenska Academy at the University of Gothenburg, Mölndal, Sweden

<sup>2</sup>Clinical Neurochemistry Laboratory, Sahlgrenska University Hospital, Mölndal, Sweden

<sup>3</sup>Neurology Clinic, University of Perugia, Perugia, Italy

<sup>4</sup>Department of Pharmaceutical Sciences, University of Perugia, Perugia, Italy

<sup>5</sup>Clinical Memory Research Unit, Department of Clinical Sciences Malmö, Lund University, Lund, Sweden

<sup>6</sup>Memory Clinic, Skåne University Hospital, Malmö, Sweden

<sup>7</sup>Department of Molecular Neuroscience, University College London Institute of Neurology, Queen Square, London, UK

<sup>8</sup>UK Dementia Research Institute at UCL, London, United Kingdom

Corresponding Author: Simon Sjödin, Department of Psychiatry and Neurochemistry, Institute of Neuroscience and Physiology, the Sahlgrenska Academy at the University of Gothenburg, House V3, SU/Mölndal, SE-43180, Mölndal, Sweden. simon.sjodin@neuro.gu.se.

## Content

Figure S4

**Figure S4. CSF protein concentrations in** *APOE*  $\varepsilon$ **4 carriers.** In Clinical Study II a subset of the participants had known *APOE* genotype. *APOE*  $\varepsilon$ **4** carriers (*APOE*  $\varepsilon$ **4** +), having one or two  $\varepsilon$ **4** alleles, were compared to non-carriers (*APOE*  $\varepsilon$ **4** –). The carriers vs. non-carriers where compared within the groups, controls (carriers, N = 10, non-carriers, N = 17) and AD (carriers, N = 25, non-carriers, N =10), respectively. Significant different CSF concentrations were identified between control carriers and control non-carriers for peptides (A) C9\_232-242 (P <0.05), (B) C9\_497-508 (P <0.05), (C) CTSF\_442-450 (P <0.05), (D) DPP7\_113-123 (P <0.05) and (E) GM2A\_89-96 (P <0.05). Statistics were calculated using Wilcoxon 2-sample rank sum test and the graphs show Tukey boxplots.





DPP7\_113-123

GM2A\_89-96



Ε