

**Table 1.** Results of urinary collection performed upon hospital admission and ten months after cessation of octreotide therapy.

<b>Parameters</b>	<b>During hospitalisation</b>	<b>10 months after hospitalization and stopping octreotide therapy</b>	<b>Normal values</b>
<b>NaFE (%)</b>	<b>10.1</b>	<b>1.07</b>	<b>&lt;1</b>
<b>UFE (%)</b>	<b>49.4</b>	<b>36.6</b>	<b>&lt;35</b>
<b>UCa (micromol/ 24h)</b>	<b>2.24</b>	<b>1.2</b>	<b>2.5-7.5</b>
<b>UPh (micromol/ 24h)</b>	<b>5.4</b>	<b>19.8</b>	<b>13-42</b>
<b>UOx (micromole/ 24h)</b>	<b>1150</b>	<b>255</b>	<b>78-489</b>
<b>UGlycerate/Cr ( mmol/mol)</b>	<b>7</b>		<b>&lt;108</b>
<b>UGlycolate/Cr (micromol/mol)</b>	<b>2</b>		<b>&lt;6</b>
<b>UCitrate/Cr (micromol/mol)</b>	<b>12</b>		<b>&lt;457</b>
<b>UIsocitrate/Cr (micromol/mol)</b>	<b>8</b>		<b>&lt;49</b>
<b>POx (micromole/l)</b>	<b>5</b>		<b>&lt;33</b>