Depression and Anxiety in Rosacea Patients: A Systematic Review and Meta-Analysis

Ru Dai, MD¹; BingJiang, Lin³; Xuetong Zhang, MD¹; Yuchen Lou, MD¹; Suling Xu, BS^{2*}

¹Department of Dermatology, Second Affiliated Hospital, School of Medicine, Zhejiang University, No. 88 Jiefang Road, Hangzhou, Zhejiang, 310009, P.R. China ²Department of Dermatology, The affiliated Hospital of Medical School of Ningbo University, No. 247, Renmin Road, Ningbo, Zhejiang, 315010, P.R. China

³ Department of Dermatology, Ningbo First Hospital, Zhejiang University, No. 59, Liuting Street, Ningbo, Zhejiang, 315010, China

Short Title: depression and anxiety in patients with rosacea


*Corresponding Author: Suling Xu, BS, Department of Dermatology, The affiliated

Hospital of Medical School of Ningbo University, No. 247, Renmin Road, Ningbo, Zhejiang,


315010, P.R. China(Tel: 86+15201017376; Fax: 86+0574-87783581; E-mail: xusuling@nbu.edu.cn)

Figure. S1. Sensitivity analysis of studies focusing on the odds ratios for depression and anxiety in rosacea patients compared with healthy controls

(a) Depression



(b) Anxiety



- (a) Sensitivity analysis of studies focusing on depression.
- (b) Sensitivity analysis of studies focusing on anxiety.

Table S1. Study quality assessment using the Newcastle-Ottawa Scale

Study	Design	Selection	Comparability	Outcome/exposure	Total
Lukaviciute et al. 2020	Case-control	3	2	1	7
Uysal et al, 2019	Case-control	2	2	3	7
Alinia et al, 2018	Cross-sectional	2	/	3	5
Huang et al, 2018	Cohort	4	2	3	9
Singam et al, 2018	Cohort	3	2	3	8
Wu et al, 2017	Cross-sectional	3	2	2	7
Egeberg et al, 2016	Cohort	4	2	3	9
Halioua et al, 2016	Cross-sectional	3	2	2	7
Dirschka et al, 2015	Cross-sectional	3	2	2	7
Böhm et al, 2014	Cross-sectional	2	0	2	4
Spoendlin et al, 2013	Case-control	3	2	3	8
Beaulieu et al, 2012	Cross-sectional	3	/	2	5
Abram et al, 2009	Case-control	4	0	2	6
Gupta et al, 2005	Cohort	3	2	3	8