Title

Maternal and Perinatal Risk Factors for Infantile Hemangioma: a Matched Case-Control Study With a Large Sample Size

Xue Gong^{1†}, MD, Tong Qiu^{1†}, MD, Liwei Feng^{1†}, BSN, Kaiying Yang¹, MD, Shiyi Dai¹, MD, Jiangyuan Zhou¹, MD, Xuepeng Zhang¹, MD, Siyuan Chen², MD, PhD, Yi Ji^{1*}, MD, PhD

¹Division of Oncology, Department of Pediatric Surgery, West China Hospital of Sichuan University, Chengdu, 610041, China

²Pediatric Intensive Care Unit, Department of Critical Care Medicine, West China Hospital of Sichuan University, Chengdu, 610041, China

†: These authors contributed equally to this work and shared first authorship.

*Address correspondence to: Yi Ji, MD, PhD, Division of Oncology, Department of Pediatric Surgery. West China Hospital of Sichuan University. #37# Guo-Xue-Xiang, Chengdu, 610041, China. Tel: +86 18980606865; Fax: +86 28 85423453; Email: jijjiyuanyuan@163.com.

Characteristics	n (%)
Female to male ratio	2.17:1
Localization	
Trunk	362(35.04%)
Face	280(27.11%)
Extremities	263(25.46%)
Head and neck	175(16.94%)
Perineum	51(4.94%)
Description	
Superficial	738(71.44%)
Fixed	220(21.30%)
Deep	46(4.45%)
Morphological subtype	
Focal	857(82.96%)
Multifocal	61(5.91%)
Indeterminate	64(6.20%)
Segmental	59(5.71%)
Complications	
Disfigurement	235(22.75%)
Ulceration	41(3.97%)
Visual impairment	8(0.77%)
Heart failure	3(0.29%)

Supplementary Table S1: Part of the Clinical Characteristics of the IH Cases