

A prospective, monocentric case-control study on uncontrolled psoriasis as independent risk factor for a hypercoagulable state

Tom M Hillary¹MD, Tine Vanhoutvin¹, Miet Peeters², Maya Imbrechts²PhD, Thomas Vanassche³MD,PhD Marjan Garmyn¹MD,PhD, Séverine Vermeire⁴ MD,PhD

¹: Dermatology department, University Hospitals Leuven, Herestraat 49, 3000, Leuven, Belgium

²: KU Leuven, Laboratory for Therapeutic and Diagnostic Antibodies, Department of Pharmaceutical and Pharmacological Sciences, Leuven, Belgium

³: Department of Cardiovascular Diseases, University Hospitals Leuven, Herestraat 49, 3000, Leuven, Belgium

⁴: Department of Gastroenterology & Hepatology, University Hospitals Leuven and Translational Research in Gastrointestinal Disorders (TARGID) KU Leuven, Belgium

Corresponding author: Tom Hillary

Herestraat 49

3000 Leuven

Belgium

+32 16 33 79 50

Tom.Hillary@uzleuven.be

Appendix A

	50%CLT_BL	50%CLT_FU	P-value
IL-17 inh	146,4	151,8	0,0509
IL-23 inh	142,6	188,6	0,0007
Systemics	150,6	168,6	0,8196
TNF inh	131,3	152,3	0,5231

Appendix A: treatment with selective IL23-antibodies was associated with a significantly prolonged 50%CLT ($p=0.0007$). This was not observed in the other treatment groups. IL: interleukin; inh: inhibitor; TNF: Tumor necrosis factor α .

Appendix B

BSA>15	mean	Q1	Q3	p
AUC % HC	213.1	134.2	281.3	
AUC % Pso	294.8	185.0	381.2	<i>0.0003</i>
Amplitude % HC	136.4	114.3	154.8	
Amplitude % Pso	161.1	144.0	177.5	<i><.0001</i>
BSA>20				
AUC % HC	207.3	140.0	269.1	
AUC % Pso	293.6	232.7	325.9	<i>0.0019</i>
Amplitude % HC	140.6	116.8	158.4	
Amplitude % Pso	165.2	142.0	178.6	<i>0.0049</i>
BSA>15				
AUC % Baseline	294.8	185.0	381.2	
AUC % Follow-up	324.9	189.2	406.8	<i>0.3743</i>
Amplitude % Baseline	161.1	144.0	177.5	
Amplitude % Follow-up	154.7	125.9	172.2	<i>0.0378</i>
BSA>20				
AUC % Baseline	293.6	232.7	325.9	
AUC % Follow-up	314.5	194.4	404.3	<i>0.8084</i>
Amplitude % Baseline	165.2	142.0	178.6	
Amplitude % Follow-up	153.9	132.9	170.8	<i>0.0084</i>

Appendix B: Increasing the threshold of BSA to 15% or 20% did not alter the effect on amplitude or AUC. HC: healthy control; Pso: psoriasis; BSA: body surface area.