SUPPLEMENTARY MATERIAL

The Epidemiology of Hospital-Treated Alopecia Areata in Denmark, 1995-2016

Sissel Brandt Toft Sørensen,¹ Prethibha George,² Oladayo Jagun,² Robert Wolk³, Lynne Napatalung^{2,4}, Samuel H. Zwillich³, Lars Iversen,^{5,6} Vera Ehrenstein¹

¹Department of Clinical Medicine and Department of Clinical Epidemiology, Aarhus University and

Aarhus University Hospital, Aarhus N, Denmark

²Pfizer Inc., New York, NY, USA

³Pfizer Inc, Groton, CT, USA

⁴Department of Dermatology, Icahn School of Medicine at Mount Sinai, New York, NY, USA

⁵Department of Dermatology, Aarhus University Hospital and Aarhus University, Aarhus N,

Denmark

⁶MC2 Therapeutics A/S, Hørsholm, Denmark

Corresponding author: Prof Vera Ehrenstein, Department of Clinical Medicine and Department of Clinical Epidemiology, Aarhus University and Aarhus University Hospital, Olof Palmes Allé 43-45 8200 Aarhus N, Denmark; Tel: +45 87 16 88 97, E-mail: ve@clin.au.dk

Table S1. ICD-10 Codes to Identify Hospital-Treated AA

Description		Severity Subtype*		
	ICD-10	AT/AU	Non-AT/AU	All AA
Alopecia areata	L63		×	×
Alopecia (capitis) totalis	L63.0	×		×
Alopecia universalis	L63.1	×		×
Ophiasis	L63.2		×	×
Other alopecia areata	L63.8		×	×
Alopecia areata, unspecified	L63.9		×	×

*For patients falling in more than one severity subtype based on the first-recorded ICD-10 codes recorded on the AA cohort entry date, patients were classified as AT/AU if they had the respective code. Otherwise they were classified as non-AT/AU to create mutually exclusive all-inclusive categories. AA, alopecia areata; AT, alopecia totalis; AU, alopecia universalis.