

| | Study participants n = 391 | Non-survivors n = 234 | Survivors n = 157 | Descriptive p-value (unadjusted) ^a |
|--|-------------------------------|--------------------------|----------------------|--|
| Co-managing specialty | | | | 0.153 |
| Visceral Surgery | 121 (31) | 69 (30) | 52 (33) | |
| Gastroenterology and Hepatology | 63 (16) | 47 (20) | 16 (10) | |
| Cardiology | 35 (9) | 21 (9) | 14 (9) | |
| Pneumology | 32 (8) | 20 (9) | 12 (8) | |
| Cardiovascular Surgery | 21 (5) | 13 (6) | 8 (5) | |
| Trauma and Orthopedic Surgery | 20 (5) | 10 (4) | 10 (6) | |
| Oncology, Hematology and Bone Marrow Transplantation | 19 (5) | 14 (6) | 5 (3) | |
| Vascular Medicine | 14 (4) | 8 (3) | 6 (4) | |
| Nephrology | 13 (3) | 5 (2) | 8 (5) | |
| Visceral Transplant Surgery | 12 (3) | 7 (3) | 5 (3) | |
| Neurology | 10 (3) | 6 (3) | 4 (3) | |
| Neurosurgery | 9 (2) | 3 (1) | 6 (4) | |
| Urology | 9 (2) | 3 (1) | 6 (4) | |
| Otorhinolaryngology | 6 (2) | 2 (1) | 4 (3) | |
| Gynecology | 4 (1) | 3 (1) | 1 (1) | |
| Dermatology | 3 (1) | 3 (1) | 0 (0) | |
| Surgical vs. medical admission | | | | 0.019 |
| Surgery | 216 (55) | 118 (50) | 98 (62) | |
| Conservative medicine | 175 (45) | 116 (50) | 59 (38) | |