

P Patients with bowel anastomosis
I Stapling with bloodclotting adjuvant
C Standard stapling
O Mortality, morbidity, anastomotic leak

P Patients with bowel anastomosis
I Stapling with 4 or more rows
C Standard stapling
O Mortality, morbidity, anastomotic leak

P Patients with bowel anastomosis
I Stapler anastomosis
C Hand-sewn anastomosis
O Mortality, morbidity, anastomotic leak

P Patients with distal pancreas resection
I Stapler closure
C Hand-sewn closure
O Mortality, morbidity, anastomotic leak

P Patients undergoing haemorrhoidectomy
I Stapler operation (Longo)
C Conventional technique
O Mortality, morbidity, re-operation, operation time

((("stapler"[tiab] OR "Surgical Staplers"[Mesh] OR "surgical stapler" [tiab] OR "surgical staplers" [tiab] OR "Surgical Stapling" [MAJR] OR "surgical stapling"[tiab] OR (stapl* AND (row OR rows) OR "stapler anastomosis" OR "stapler anastomoses" OR "stapler closure" Or "milligan-morgan")) AND (surgery[tiab] OR surgeries[tiab] OR surgic*[tiab])) AND ((randomized controlled trial[pt] OR controlled clinical trial[pt] OR randomized[tiab] OR placebo[tiab] OR clinical trials as topic[mesh:noexp] OR double blind method[mh] OR randomly[tiab] OR trial[ti] NOT (animals[mh] NOT humans[mh])))