

Table A1. Missing values for health indicators by survey design and mode of data collection

	Concurrent mixed-mode design			Sequential mixed-mode design		
	SAQ-Web (n=124)	SAQ-Paper (n=482)	CATI (n=11)	SAQ-Web (n=290)	SAQ-Paper (n=264)	CATI (n=7)
	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)
Diabetes (12-month prevalence)	0 (0.0)	36 (7.5)	0 (0.0)	0 (0.0)	18 (6.8)	0 (0.0)
Hypertension (12-month prevalence)	0 (0.0)	40 (8.3)	0 (0.0)	0 (0.0)	18 (6.8)	0 (0.0)
Coronary heart disease (lifetime prevalence)	0 (0.0)	7 (1.5)	0 (0.0)	0 (0.0)	4 (1.5)	0 (0.0)
Chronic bronchitis (lifetime prevalence)	1 (0.8)	35 (7.3)	1 (9.1)	0 (0.0)	17 (6.4)	0 (0.0)
Obesity (BMI \geq30)	0 (0.0)	10 (2.1)	0 (0.0)	6 (2.1)	12 (4.6)	0 (0.0)
Current depression	11 (8.9)	36 (7.5)	0 (0.0)	7 (2.4)	31 (11.7)	1 (14.3)
Mental well-being	4 (3.2)	26 (5.4)	0 (0.0)	10 (3.5)	22 (8.3)	0 (0.0)
Social support	0 (0.0)	9 (1.9)	0 (0.0)	0 (0.0)	9 (3.4)	1 (14.4)
Current smoking	0 (0.0)	13 (2.7)	0 (0.0)	0 (0.0)	11 (4.2)	0 (0.0)
At-risk alcohol consumption	2 (1.6)	51 (10.6)	0 (0.0)	6 (2.1)	33 (12.5)	0 (0.0)
Sport participation	0 (0.0)	13 (2.7)	0 (0.0)	0 (0.0)	3 (1.1)	0 (0.0)
Influenza vaccination (2011/2012)	0 (0.0)	20 (4.2)	0 (0.0)	1 (0.3)	9 (3.4)	0 (0.0)

SAQ-Web = self-administered web survey; SAQ-Paper = self-administered paper mail questionnaire; CATI = computer-assisted telephone interview.