
TAIC-I Trial Protocol
Administration of TAICs
Immunosuppression with Tacrolimus, Sirolimus and corticosteroids
Steroid weaning phase
Sirolimus weaning phase
Minimisation of Tacrolimus treatment (8-10ng/ml)
Further reduction of Tacrolimus therapy
Follow-up
WW
Excluded
FK
Excluded
DW
Exit: Day 28, DVT
UF
Exit: Day 39, Rejection – Banff Grade III
MR
Exit: Day 53, Unconfirmed rejection episode
ME
Exit: Day 54, Unconfirmed rejection episode
MT
Exit: Day 58, Rejection – Banff Grade II
GH
Exit: Day 63, Rejection – Banff Grade I
KGW
Exit: Day 84, Unconfirmed rejection episode
UH
Exit: Day 164, Withdrew consent
GS
Exit: Day 168, Withdrew consent
GM
Exit: Day 193, Rejection – Banff Grade I
0
4
8
12
16
20
24
28
32
Time post-transplantation (weeks)
Legend
Exit from the trial with no evidence of rejection
Exit from the trial with a clinical suspicion of rejection, but without histological confirmation
Exit from trial with histologically confirmed rejection
Complete withdrawal of all immunosuppressive therapy
Tacrolimus monotherapy, trough serum levels: ≤4.1ng/ml
Tacrolimus monotherapy, trough serum levels: ≥5 - 8 ng/ml
Tacrolimus monotherapy, trough serum levels: ≥8 - 10 ng/ml