

**Burden of first osteoporotic hip fracture in Spain: a prospective, observational, 12-month study**

*Calcified Tissue International*

**Authors:** Jose Ramón Caeiro (1), Agustí Bartra (2), Manuel Mesa-Ramos (3), Íñigo Etxebarria (4), Jorge Montejo (5), Pedro Carpintero (6), Francesc Sorio (7), Sonia Gatell (7), Andrea Farré (7), and Laura Canals (7) on behalf of the PROA investigators

**Corresponding author:**

Dr. José Ramón Caeiro

Servicio de COT

Complejo Hospitalario de Santiago de Compostela

Calle Choupana s/n, ES-15702 Santiago de Compostela, A Coruña, Spain.

Phone: +34-609458500

Fax: +34-981950900

Email: jrcaeiro@telefonica.net

**Supplementary Table 1.** Changes over time in living arrangements

	Baseline Living Arrangement					
	Own home alone	Own home with partner / family member	Nursing home	Relatives home	Unknown	Total
	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)
<b>4 months</b>						
Own home alone	37 (7.6)	13 (2.7)	2 (0.4)	2 (0.4)	-	54 (11.1)
Own home with partner / family member	29 (6.0)	200 (41.1)	3 (0.6)	12 (2.5)	-	244 (50.1)
Nursing home	12 (2.5)	18 (3.7)	36 (7.4)	4 (0.8)	-	70 (14.4)
Relatives home	4 (0.8)	17 (3.5)	0 (0.0)	14 (2.9)	-	35 (7.2)
Unknown	11 (2.2)	50 (10.3)	14 (2.9)	8 (16.4)	1 (0.2)	84 (17.2)
Total	93 (19.1)	298 (61.2)	55 (11.3)	40 (8.2)	1 (0.2)	487 (100)
<b>12 months</b>						

Own home alone	36 (7.4)	15 (3.1)	2 (0.4)	1 (0.2)	-	54 (11.1)
Own home with partner / family member	28 (5.7)	183 (37.6)	5 (1.0)	12 (2.5)	-	228 (46.8)
Nursing home	10 (2.1)	15 (3.1)	26 (5.3)	5 (1.0)	-	56 (11.5)
Relatives home	3 (0.6)	15 (3.1)	0 (0.0)	9 (1.8)	-	27 (5.5)
Unknown	16 (3.3)	70 (14.4)	22 (4.5)	13 (2.7)	1 (0.2)	122 (25.1)
Total	93 (19.1)	298 (61.2)	55 (11.3)	40 (8.2)	1 (0.2)	487 (100)

Percentages based on total number of patients (N=487)