

**Psychometric validation of the Spanish version of the Expanded Prostate cancer
Index Composite-26**

World Journal of Urology

Víctor Zamora MPH^{1,2,3}, Olatz Garin PhD^{1,3,4*}, José Francisco Suárez MD⁵, Josep Jové MD⁶, Manuel Castells MD⁵, Ferran Ferrer MD⁷, Cristina Gutiérrez PhD⁷, Ferran Guedea PhD⁷, Ana Boladeras MD⁷, Lluís Fumadó PhD⁸, Alvar Roselló MD⁹, Jorge Pastor MD¹⁰, Pilar Samper PhD¹¹, Àngels Pont MSc^{1,3}, Montse Ferrer PhD^{1,3,4*}

1 Health Services Research Group, Hospital del Mar Research Institute, Barcelona, Spain.

2 Universitat Autònoma de Barcelona (UAB), Bellaterra, Spain.

3 CIBER en Epidemiología y Salud Pública, CIBERESP, Spain.

4 Universitat Pompeu Fabra, Barcelona, Spain.

5 Urology Department, Hospital Universitari de Bellvitge, L'Hospitalet de Llobregat, Spain.

6 Radiation Oncology Department, Institut Català d'Oncologia, Badalona, Spain.

7 Radiation Oncology Department, Institut Català d'Oncologia, L'Hospitalet de Llobregat, Spain.

8 Urology Department, Hospital del Mar, Barcelona, Spain.

9 Radiation Oncology Department, Institut Català d'Oncologia, Girona, Spain.

10 Radiation Oncology Department, ASCIRES GRUPO BIOMÉDICO, Valencia, Spain.

11 Radiation Oncology Department, Hospital Universitario Rey Juan Carlos, Móstoles, Spain

Contact information for corresponding authors

*Olatz Garin and Montse Ferrer, PhD,

e-mail address: ogarin@imim.es; mferrer@imim.es

Supplementary Table 1. Absolute and relative frequencies of EPIC-26 response options at 12 months after treatment (n=543).

EPIC-26		Response options, n (%)				
Items, n (%)	The Best				The Worst	
Urinary incontinence						
1. Leaking	311 (57.3%)	95 (17.5%)	54 (9.9%)	31 (5.7%)	52 (9.6%)	
2. Dribbling	295 (54.3%)	148 (27.3%)	95 (17.5%)	5 (0.9%)	---	
3. Pad use*	455 (83.8%)	57 (10.5%)	20 (3.7%)	11 (2.0%)	---	
4a. Leaking problem*	294 (54.1%)	5 (0.9%)	57 (10.5%)	95 (17.5%)	92 (16.9%)	
Urinary Irritative/Obstructive						
4b. Dysuria*	462 (85.1%)	1 (0.2%)	32 (5.9%)	30 (5.5%)	18 (3.3%)	
4c. Hematuria*	535 (98.5%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	8 (1.5%)	
4d. Weak stream*	362 (66.7%)	2 (0.4%)	43 (7.9%)	70 (12.9%)	66 (12.2%)	
4e. Frequency*	267 (49.2%)	1 (0.2%)	53 (9.8%)	127 (23.4%)	95 (17.5%)	
5. Overall urinary problem*	194 (35.7%)	1 (0.2%)	87 (16.0%)	148 (27.3%)	113 (20.8%)	
Bowel						
6a. Urgency*	486 (89.5%)	0 (0.0%)	11 (2.0%)	27 (5.0%)	19 (3.5%)	
6b. Frequency*	462 (85.1%)	0 (0.0%)	13 (2.4%)	42 (7.7%)	26 (4.8%)	
6c. Fecal incontinence*	480 (88.4%)	0 (0.0%)	2 (0.4%)	20 (3.7%)	41 (7.6%)	
6d. Bloody stools*	492 (90.6%)	0 (0.0%)	15 (2.8%)	19 (3.5%)	17 (3.1%)	
6e. Rectal pain*	474 (87.3%)	0 (0.0%)	13 (2.4%)	39 (7.2%)	17 (3.1%)	
7. Overall bowel problem*	372 (68.5%)	0 (0.0%)	40 (7.4%)	79 (14.5%)	52 (9.6%)	
Sexual						
8a. Poor erections	0 (0.0%)	99 (18.2%)	59 (10.9%)	83 (15.3%)	302 (55.6%)	
8b. Difficulty with orgasm	2 (0.4%)	128 (23.6%)	74 (13.6%)	44 (8.1%)	295 (54.3%)	
9. Erection not firm	123 (22.7%)	87 (16.0%)	64 (11.8%)	269 (49.5%)	---	
10. Erections frequency	86 (15.8%)	47 (8.7%)	61 (11.2%)	24 (4.4%)	325 (59.9%)	
11. Poor sexual function	0 (0.0%)	89 (16.4%)	63 (11.6%)	58 (10.7%)	333 (61.3%)	
12. Overall sexual problem*	355 (65.4%)	0 (0.0%)	31 (5.7%)	86 (15.8%)	71 (13.1%)	
Hormonal						
13a. Hot flashes*	384 (70.7%)	0 (0.0%)	20 (3.7%)	46 (8.5%)	93 (17.1%)	
13b. Breast problems*	437 (80.5%)	4 (0.7%)	54 (9.9%)	38 (7.0%)	10 (1.8%)	
13c. Depression*	312 (57.5%)	1 (0.2%)	29 (5.3%)	82 (15.1%)	119 (21.9%)	
13d. Lack of energy*	307 (56.5%)	0 (0.0%)	25 (4.6%)	78 (14.4%)	133 (24.5%)	
13e. Weight change*	415 (76.4%)	2 (0.4%)	31 (5.7%)	51 (9.4%)	44 (8.1%)	

**The direction of these items is opposite (lower scores mean better outcomes)*