

Study or Subgroup	Laser Haemorrhoidoplasty			Milligan-Morgan (Open) Haemorrhoidectomy			Mean Difference	
	Mean [min]	SD [min]	Total	Mean [min]	SD [min]	Total	Weight	IV, Random, 95% CI
Nazari 2015	32.8	7	29	52.3	15.1	30	8.3%	-19.50 [-25.47, -13.53]
Naderan 2016	33.1	7.3	30	52.6	15.6	30	8.1%	-19.50 [-25.66, -13.34]
Alsisy 2019	30.63	4.9	30	50.5	10.12	30	10.6%	-19.87 [-23.89, -15.85]
Eskandaros 2020	22.8	3.9	40	27.5	5.3	40	12.9%	-4.70 [-6.74, -2.66]
Poskus 2020	15	7.7	40	29	13.8	40	9.5%	-14.00 [-18.90, -9.10]
Maloku 2014	15.94	3.5	20	26.76	5.8	20	11.9%	-10.82 [-13.79, -7.85]
Maloku 2019	16.5	1.9	100	27.2	6.5	100	13.5%	-10.70 [-12.03, -9.37]
Hassan 2021	15.9	3.5	20	26.8	5.8	20	11.9%	-10.90 [-13.87, -7.93]
Khan 2021	23.2	1.9	50	34.3	5.5	50	13.3%	-11.10 [-12.71, -9.49]
Total (95% CI)			359			360	100.0%	-12.74 [-15.43, -10.05]

Heterogeneity: Tau² = 13.50; Chi² = 71.37, df = 8 (P < 0.00001); I² = 89%
 Test for overall effect: Z = 9.29 (P < 0.00001)

