

Moderate consumption of fermented alcoholic beverages diminishes diet-induced non-alcoholic fatty liver disease through mechanisms involving hepatic adiponectin signaling in mice

Finn Jung, Tino Lippmann, Annette Brandt, Cheng Jun Jin, Anna Janina Engstler, Anja Baumann^{*}

Article type: Original article

European Journal of Nutrition

*** Corresponding author:** Anja Baumann, Ph.D.
University of Vienna
Department of Nutritional Sciences
Molecular Nutritional Science
Althanstraße 14 (UZA II)
A-1090 Wien
Email address: anja.baumann@univie.ac.at

Online Resource 3. Primer sequences used for real-time RT-PCR.

Gene	Forward 5'---3'	Reverse 5'---3'
<i>18s</i>	ccatccaatcggtagtagcg	gtaaccegtgaacccatt
<i>Acox1</i>	gggaattggcatcgaga	catctccgtctgggcgtagg
<i>Adipoq</i>	gaagacctgcatctcctttctctcc	tcatgccgaagatgacgttactaca
<i>Adipor1</i>	gggacttggcttgagtggg	gggaagaagcgtcaggaat
<i>Adipor2</i>	tgccttaggtaggccaaca	tccaaagtgcacgaccaga
<i>Fas</i>	tctggccaacctcattggt	gaagctgggggtccattgtg
<i>Foxo1</i>	ctcccggtacttctctgctg	gtggccagttggactggtt
<i>Insr</i>	catcccgaagcgaagatcc	gagtctgattgcatgcctgcag
<i>Irs1</i>	gctctagtgttccgtgtcc	gttcccaccctagacaaaa
<i>Irs2</i>	gaagcggctaagtctcatgg	gacgggtgggtgtagaggaaa
<i>Pparγ1</i>	aacgtgaagcccatcgagga	ctgcacgtgctctgtgacga
<i>Sirt1</i>	gacgctgtggcagattgta	gaaatcccacaggagacaga
<i>Srebp-1c</i>	accggctactgctggactgc	agagcaagagggtgcatcg