

## Electronic Supplementary Material

Prebiotic effect of inulin-type fructans on fecal microbiota and short-chain fatty acids in type 2 diabetes: A randomized controlled trial


European Journal of Nutrition

Eline Birkeland<sup>1,2</sup>, Sedegheh Gharagozlian<sup>1</sup>, Kåre I. Birkeland<sup>2,3</sup>, Jørgen Valeur<sup>4,5</sup>, Ingrid Måge<sup>6</sup>, Ida Rud<sup>6</sup>, Anne-Marie Aas<sup>1,2</sup>

Ida Rud and Anne-Marie Aas share last authorship

<sup>1</sup>Section of Nutrition and Dietetics, Department of Clinical Service, Division of Medicine, Oslo University Hospital, Norway, <sup>2</sup>Institute of Clinical Medicine, University of Oslo, Norway <sup>3</sup>Department of Transplantation Medicine, Oslo University Hospital, Norway, <sup>4</sup>Department of Gastroenterology, Oslo University Hospital, Oslo, Norway, <sup>5</sup>Unger-Vetlesen Institute, Lovisenberg Diaconal Hospital, Oslo, Norway, <sup>6</sup>Nofima - Norwegian Institute of Food, Fisheries and Aquaculture Research, Ås, Norway.

Corresponding author: Eline Birkeland, eline.birkeland@ous-hf.no



**Online Resource 1.** Flow chart showing all patients approached for the study. \*n=12 for microbiota analysis