

Clinical Pharmacokinetics

Pipamperone population pharmacokinetics related to effectiveness and side-effects in children and adolescents

Sanne M. Kloosterboer^{1,2}, Karin M. Egberts³, Brenda C.M. de Winter¹, Teun van Gelder¹, Manfred Gerlach³, Manon H.J. Hillegers², Gwen C. Dieleman², Soma Bahmany¹, Catrien G. Reichart⁴, Emma van Daalen⁵, Mirjam E.J. Kouijzer⁶, Bram Dierckx², Birgit C.P. Koch¹

¹Department of Hospital Pharmacy, Erasmus MC, University Medical Center Rotterdam, Rotterdam, The Netherlands

²Department of Child and Adolescent Psychiatry/Psychology, Erasmus MC, University Medical Center Rotterdam, Rotterdam, the Netherlands

³ Department of Child and Adolescent Psychiatry, Psychosomatics and Psychotherapy, University Hospital Würzburg, Center of Mental Health, Würzburg, Germany

⁴ Curium-LUMC Child and Adolescent Psychiatry, Leiden University Medical Center, Leiden, The Netherlands

⁵ Yulius Mental Health, Dordrecht, The Netherlands

⁶ GGz Breburg, Centre of Youth, Breda, the Netherlands

Corresponding author:

Sanne Maartje Kloosterboer

Department of Hospital Pharmacy, Erasmus MC, University Medical Center Rotterdam

Dr.Molewaterplein 40

3015 GD Rotterdam, The Netherlands

s.kloosterboer@erasmusmc.nl

Figure S1: Goodness Of Fit plots – external validation

Figure S1a: External validation: measured concentrations versus population predictions.




Figure S1b: External validation: measured concentrations versus individual predictions.




Figure S1c: External validation: the correlation of conditional weighted residuals (CWRES) with time




Figure S1d: External validation: the correlation of conditional weighted residuals with population predicted concentrations




Figure S2: Normalized prediction distribution errors (npde) of external validation

