

SUPPLEMENTARY FIGURES FOR

“Sex-specific developmental models for *Creophilus maxillosus* (L.) (Coleoptera: Staphylinidae): searching for larger accuracy of insect age estimates”

International Journal of Legal Medicine

Katarzyna Frątczak-Łagiewska M.Sc.^{1,2*}, Szymon Matuszewski Ph.D.¹

¹Laboratory of Criminalistics, Adam Mickiewicz University, Św. Marcin 90, 61-809 Poznań, Poland.


²Department of Animal Taxonomy and Ecology, Adam Mickiewicz University, Umultowska 89, 61-614 Poznań, Poland

*corresponding author, katarzyna.fratzczak@amu.edu.pl


Content



Fig. 1. Differences between males and females of *C. maxillosus* in duration of life stages at 15°C, 17.5°C, 20°C, 22.5°C, 25°C, 27.5°C, 30°C; ■ – mean, whiskers – standard error of the mean; E – the egg stage, L1 – first larval stage, L2 – second larval stage, L3 – third larval stage, P – pupal stage.

15°C




17.5°C






25°C



27.5°C






Fig. 1. Differences between males and females of *C. maxillosus* in duration of life stages at 15°C, 17.5°C, 20°C, 22.5°C, 25°C, 27.5°C, 30°C; ■ – mean, whiskers – standard error of the mean; E – the egg stage, L1 – first larval stage, L2 – second larval stage, L3 – third larval stage, P – pupal stage.