

Article title: Prediction of renal outcome in Henoch-Schönlein nephritis based on biopsy findings

Journal: Pediatric Nephrology

Author names: Mikael Koskela, Elisa Ylinen, Helena Autio-Harmainen, Heikki Tokola, Päivi Heikkilä, Jouko Lohi, Hannu Jalanko, Matti Nuutinen, Timo Jahnukainen

Corresponding author: Mikael Koskela; Children's Hospital, Pediatric Research Center, University of Helsinki, Helsinki University Hospital, Helsinki, Finland; e-mail address: mikael.koskela@helsinki.fi

Table S1. Histologic parameters of the semiquantitative classification (SQC)

	Active or chronic	Scoring
Glomerular parameters		
Lobulation	active	0-1 ^a
Mesangial proliferation	active	0-1 ^a
Crescents		
cellular	active	0-3 ^b
fibrous	chronic	0-3 ^b
Adhesions	chronic	0-3 ^b
Fibrinous thrombosis	active	0-3 ^b
Global sclerosis	chronic	0-3 ^b
Segmental sclerosis	chronic	0-2 ^c
Tubular parameters		
Thickening of the basement membrane	chronic	0-1 ^a
Complete atrophy	chronic	0-1 ^a
Tubular dilatation	active	0-1 ^a
Interstitial parameters		
Fibrosis	chronic	0-1 ^a
Inflammation OR periglomerular inflammation	chronic	0-1 ^a
Capillary parameters		
Arteriosclerosis OR arterial inflammation	chronic	0-1 ^a
Focal or diffuse mesangial proliferation		0 for focal, 1 for diffuse

^a0 = absent; 1 = present

^b0 = 0% of glomeruli affected; 1 = 0-5% of glomeruli affected; 2 = 5-10% of glomeruli affected; 3 = >10% of glomeruli affected

^c0 = 0% of glomeruli affected; 1 = 0-5% of glomeruli affected; 3 = >5% of glomeruli affected

SQC was originally published in: The ISKDC classification and a new semiquantitative classification for predicting outcomes of Henoch-Schonlein purpura nephritis. *Pediatr Nephrol* 2017; 32: 1201-1209.