

Online Resource 4

The palatability of oral nutritional supplements: before, during, and after chemotherapy

Irene IJpma, Remco J. Renken, Gert J. ter Horst, Anna K. L. Reyners*

*Top Institute Food and Nutrition, Wageningen, The Netherlands; Department of Medical Oncology, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands. Email: a.k.l.reyners@umcg.nl.

Table I Median (IQR) difference in liking compared to baseline and per cycle among patients.

	T1 – T0 (N = 18)	T2 – T0 (N = 18)	T3 – T0 (N = 16)	T4 – T0 (N = 17)	T3-T1 (N = 15)	T3-T2 (N = 16)
Milk-based Vanilla	-0.50 (-1.25 – 0.25)	0.00 (-2.00 – 0.00)	0.00 (-2.00 – 0.00)	0.00 (-1.50 – 0.00)	0.00 (-1.00 – 0.00)	0.00 (-1.00 – 0.00)
Milk-based Strawberry	0.00 (0 – 1.00)	-0.50 (-1.25 – 1.00)	0.00 (-1.75 – 0.75)	0.00 (-1.50 – 1.00)	-1.00 (-2.00 – 0.00)	0.00 (-1.00 – 1.00)
Juice-based Apple	-1.00 (-2.00 – 1.00)	0.00 (-1.00 – 1.00)	0.50 (-1.00 – 1.00)	0.00 (-0.50 – 1.00)	0.00 (0.00 – 1.00)	0.00 (-0.75 – 1.00)
Juice-based Orange	0.00 (-2.00 – 1.00)	0.00 (-1.25 – 1.25)	0.50 (-2.00 – 2.00)	1.00 (-0.50 – 1.00)	0.00 (-1.00 – 2.00)	0.00 (0.00 – 1.00)
Yoghurt-based Vanilla-Lemon	0.00 (-1.25 – 1.00)	0.00 (-2.00 – 1.25)	0.00 (-1.00 – 1.00)	-1.00 (-1.00 – 0.00)	0.00 (-1.00 – 1.00)	0.00 (-0.75 – 0.00)
Yoghurt-based Peach-Orange	-1.00 (-2.00 – 0.25)	0.00 (-1.25 – 1.00)	0.00 (-1.00 – 2.00)	-1.00 (-2.00 – 0.50)	0.00 (-1.00 – 2.00)	0.00 (-0.75 – 1.00)